


Problem Set 4

Model Answers

Problem 1

(a) The finite automaton N is



(b) First, we put the expression of N in Forlan's syntax

```

{states} A, B, C {start state} A {accepting states} A, B, C
{transitions}
A, 0 -> B; A, 1 -> A;
B, 0 -> B; B, 1 -> C;
C, 1 -> A
  
```

in the file `ps4-p1-fa` (see the course website), and load this file into Forlan, calling the result `fa`:

```

- val fa = FA.input "ps4-p1-fa";
val fa = - : fa
  
```

Next we load the file `ps4-p1.sml`

```

(* val inX : str -> bool

   tests whether a string over the alphabet {0, 1} is in X *)

fun inX x =
  Set.all
    (fn y => not(Str.equal(y, Str.fromString "010")))
    (StrSet.substrings x);

(* val upto : int -> str set

   if n >= 0, then upto n returns all strings over alphabet {0, 1} of
   length no more than n *)

fun upto 0 : str set = Set.sing nil
  
```

```

| upto n          =
  let val xs = upto(n - 1)
      val ys = Set.filter (fn x => length x = n - 1) xs
  in StrSet.union
    (xs, StrSet.concat(StrSet.fromString "0, 1", ys))
  end;

(* val partition : int -> str set * str set

   if n >= 0, then partition n returns (xs, ys) where:

   xs is all elements of upto n that are in X; and

   ys is all elements of upto n that are not in X *)

fun partition n = Set.partition inX (upto n);

(* val test = fn : int -> dfa -> str option * str option

   if n >= 0, then test n returns a function f such that, for all FAs
   fa, f fa returns a pair (xOpt, yOpt) such that:

   If there is an element of {0, 1}* of length no more than n that
   is in X but is not accepted by fa, then xOpt = SOME x for some
   such x; otherwise, xOpt = NONE.

   If there is an element of {0, 1}* of length no more than n that
   is not in X but is accepted by fa, then yOpt = SOME y for some
   such y; otherwise, yOpt = NONE. *)

fun test n =
  let val (goods, bads) = partition n
      in fn fa =>
          let val accepted      = FA.accepted fa
              val goodNotAccOpt = Set.position (not o accepted) goods
              val badAccOpt     = Set.position accepted bads
          in ((case goodNotAccOpt of
                NONE    => NONE
                | SOME i => SOME(ListAux.sub(Set.toList goods, i))),
             (case badAccOpt of
                NONE    => NONE
                | SOME i => SOME(ListAux.sub(Set.toList bads, i))))
          end
        end;
end;

```

(see the course website) defining the function `test` into Forlan:

```

- use "ps4-p1.sml";
[opening ps4-p1.sml]

```

```

val inX = fn : str -> bool
val upto = fn : int -> str set
val partition = fn : int -> str set * str set
val test = fn : int -> fa -> str option * str option
val it = () : unit

```

Finally, we apply `test` to arguments `10` and `fa`:

```

- test 10 fa;
val it = (NONE,NONE) : str option * str option

```

Problem 2

(a) First, we load the file `ps4-p2-fa` (see the course website) containing the expression

```

{states} A, B, C, D {start state} A {accepting states} B, C, D
{transitions}
A, % -> B | C | D;
B, 0 -> C;
C, 0 -> D; C, 1 -> B;
D, 1 -> C

```

of M in Forlan's syntax into Forlan, calling the result `fa`:

```

- val fa = FA.input "ps4-p2-fa";
val fa = - : fa

```

Next, we define a function `accPr` that finds and prints a labeled path in `fa` explaining why a Forlan string expressed as an SML string is accepted:

```

- fun accPr s =
=      LP.output("", FA.findAcceptingLP fa (Str.fromString s));
val accPr = fn : string -> unit

```

Finally, we use this function to find and display the required labeled paths:

```

- accPr "0010110";
A, % => B, 0 => C, 0 => D, 1 => C, 0 => D, 1 => C, 1 => B, 0 => C
val it = () : unit
- accPr "1001101";
A, % => C, 1 => B, 0 => C, 0 => D, 1 => C, 1 => B, 0 => C, 1 => B
val it = () : unit
- accPr "1011001";
A, % => D, 1 => C, 0 => D, 1 => C, 1 => B, 0 => C, 0 => D, 1 => C
val it = () : unit

```

(b) Continuing our Forlan session, we first load the file `ps4-p2.sml`

```

fun accLen n =
  Set.filter
    (FA.accepted fa)
    (StrSet.power(StrSet.fromString "0,1", n));

```

(see the course website) defining the function `accLen` into Forlan:

```
- use "ps4-p2.sml";
[opening ps4-p2.sml]
val accLen = fn : int -> str set
val it = () : unit
```

Then we apply it to 10, calling the resulting set of labeled paths `lps`, compute the size of `lps`, and display its elements:

```
- val lps = accLen 10;
val lps = - : str set
- Set.size lps;
val it = 94 : int
- StrSet.output("", lps);
0010101010, 0010101011, 0010101100, 0010101101, 0010110010, 0010110011,
0010110100, 0010110101, 0011001010, 0011001011, 0011001100, 0011001101,
0011010010, 0011010011, 0011010100, 0011010101, 0100101010, 0100101011,
0100101100, 0100101101, 0100110010, 0100110011, 0100110100, 0100110101,
0101001010, 0101001011, 0101001100, 0101001101, 0101010010, 0101010011,
0101010100, 0101010101, 0101010110, 0101011001, 0101011010, 0101100101,
0101100110, 0101101001, 0101101010, 0110010101, 0110010110, 0110011001,
0110011010, 0110100101, 0110100110, 0110101001, 0110101010, 1001010101,
1001010110, 1001011001, 1001011010, 1001100101, 1001100110, 1001101001,
1001101010, 1010010101, 1010010110, 1010011001, 1010011010, 1010100101,
1010100110, 1010101001, 1010101010, 1010101011, 1010101100, 1010101101,
1010110010, 1010110011, 1010110100, 1010110101, 1011001010, 1011001011,
1011001100, 1011001101, 1011010010, 1011010011, 1011010100, 1011010101,
1100101010, 1100101011, 1100101100, 1100101101, 1100110010, 1100110011,
1100110100, 1100110101, 1101001010, 1101001011, 1101001100, 1101001101,
1101010010, 1101010011, 1101010100, 1101010101
val it = () : unit
```

Problem 3

Define a function `dsfxs` (for “diffs of suffixes”) from $\{0, 1\}^*$ to $\mathcal{P}\mathbb{Z}$ by: for all $w \in \{0, 1\}^*$,

$$\mathbf{dsfxs} w = \{ \mathbf{diff} v \mid v \text{ is a suffix of } w \}.$$

From the definitions of X and `dsfxs` and the fact that suffixes are substrings, we have that, if $w \in X$, then $\mathbf{dsfxs} w \subseteq \{-2, -1, 0, 1, 2\}$. It turns out, though, that we can characterize membership in X using `dsfxs`.

Lemma PS4.3.1

For all $w \in X$ and $n, m \in \mathbf{dsfxs} w$, $-2 \leq m - n \leq 2$.

Proof. Suppose $w \in X$ and $n, m \in \mathbf{dsfxs} w$, so that there are suffixes u and v of w such that $n = \mathbf{diff} u$ and $m = \mathbf{diff} v$. Because u and v are suffixes of w , one must be a suffix of the other, and so there are two cases to consider.

- Suppose u is a suffix of v . Thus $v = zu$ for some $z \in \{0, 1\}^*$, and thus z is a substring of w . Hence $m = \mathbf{diff} v = \mathbf{diff} z + \mathbf{diff} u = \mathbf{diff} z + n$, so that $m - n = \mathbf{diff} z$. Because z is a substring of $w \in X$, we have that $-2 \leq \mathbf{diff} z \leq 2$, and thus $-2 \leq m - n \leq 2$.
- Suppose v is a suffix of u . Thus $u = zv$ for some $z \in \{0, 1\}^*$, and thus z is a substring of w . Hence $n = \mathbf{diff} u = \mathbf{diff} z + \mathbf{diff} v = \mathbf{diff} z + m$, so that $n - m = \mathbf{diff} z$. Because z is a substring of $w \in X$, we have that $-2 \leq \mathbf{diff} z \leq 2$, and thus $-2 \leq n - m \leq 2$. Since $-2 \leq n - m$, we have that $m - n = -(n - m) \leq -(-2) = 2$. And since $n - m \leq 2$, we have that $-2 \leq -(n - m) = m - n$. Thus $-2 \leq m - n \leq 2$.

□

Lemma PS4.3.2

For all $w \in X$, either $\mathbf{dsfxs} w \subseteq \{-2, -1, 0\}$ or $\mathbf{dsfxs} w \subseteq \{-1, 0, 1\}$ or $\mathbf{dsfxs} w \subseteq \{0, 1, 2\}$.

Proof. Suppose $w \in X$. Thus $\mathbf{dsfxs} w \subseteq \{-2, -1, 0, 1, 2\}$. Because $\%$ is a suffix of w , we have that $0 = \mathbf{diff} \% \in \mathbf{dsfxs} w$. There are two cases to consider.

- Suppose $-2 \in \mathbf{dsfxs} w$. Lemma PS4.3.1 tells us that neither 1 nor 2 are elements of $\mathbf{dsfxs} w$, since $-2 - 1 = -3$ and $-2 - 2 = -4$ are both < -2 . Thus $\mathbf{dsfxs} w \subseteq \{-2, -1, 0\}$, so that either $\mathbf{dsfxs} w \subseteq \{-2, -1, 0\}$ or $\mathbf{dsfxs} w \subseteq \{-1, 0, 1\}$ or $\mathbf{dsfxs} w \subseteq \{0, 1, 2\}$.
- Suppose $-2 \notin \mathbf{dsfxs} w$. Then $\mathbf{dsfxs} w \subseteq \{-1, 0, 1, 2\}$. There are two subcases to consider.
 - Suppose $2 \in \mathbf{dsfxs} w$. Then Lemma PS4.3.1 tells us that -1 is not an element of $\mathbf{dsfxs} w$, since $2 - (-1) = 3$ is > 2 . Thus $\mathbf{dsfxs} w \subseteq \{0, 1, 2\}$, so that either $\mathbf{dsfxs} w \subseteq \{-2, -1, 0\}$ or $\mathbf{dsfxs} w \subseteq \{-1, 0, 1\}$ or $\mathbf{dsfxs} w \subseteq \{0, 1, 2\}$.
 - Suppose $2 \notin \mathbf{dsfxs} w$. Then $\mathbf{dsfxs} w \subseteq \{-1, 0, 1\}$, so that either $\mathbf{dsfxs} w \subseteq \{-2, -1, 0\}$ or $\mathbf{dsfxs} w \subseteq \{-1, 0, 1\}$ or $\mathbf{dsfxs} w \subseteq \{0, 1, 2\}$.

□

Lemma PS4.3.3

For all $w \in \{0, 1\}^*$ and $n \in \{-2, -1, 0\}$, if $\mathbf{dsfxs} w \subseteq \{n, n + 1, n + 2\}$, then $w \in X$.

Proof. Suppose $w \in \{0, 1\}^*$, $n \in \{-2, -1, 0\}$ and $\mathbf{dsfxs} w \subseteq \{n, n + 1, n + 2\}$. To show that $w \in X$, suppose v is a substring of w . Thus $w = xvy$ for some $x, y \in \{0, 1\}^*$. We must show that $-2 \leq \mathbf{diff} v \leq 2$. Because y is a suffix of w , $\mathbf{diff} y \in \mathbf{dsfxs} w$, and thus $n \leq \mathbf{diff} y \leq n + 2$. Because vy is a suffix of w , $\mathbf{diff}(vy) \in \mathbf{dsfxs} w$, and thus $n \leq \mathbf{diff}(vy) \leq n + 2$. And since $\mathbf{diff}(vy) = \mathbf{diff} v + \mathbf{diff} y = \mathbf{diff} y + \mathbf{diff} v$, it follows that $n \leq \mathbf{diff} y + \mathbf{diff} v \leq n + 2$.

Suppose, toward a contradiction, that $-2 \leq \mathbf{diff} v \leq 2$ is false. Thus there are two cases to consider.

- Suppose $\mathbf{diff} v \leq -3$. Because $\mathbf{diff} y \leq n + 2$, it follows that $n \leq \mathbf{diff} y + \mathbf{diff} v \leq (n + 2) + (-3) = n - 1$, so that $n \leq n - 1$ —contradiction.
- Suppose $3 \leq \mathbf{diff} v$. Because $n \leq \mathbf{diff} y$, it follows that $n + 3 \leq \mathbf{diff} y + \mathbf{diff} v \leq n + 2$, so that $3 \leq 2$ —contradiction.

Because we obtained a contradiction in both cases, we have an overall contradiction. Thus $-2 \leq \mathbf{diff} v \leq 2$, completing the proof that $w \in X$. \square

Lemma PS4.3.4

For all $w \in \{0, 1\}^*$, if either $\mathbf{dsfxs} w \subseteq \{-2, -1, 0\}$ or $\mathbf{dsfxs} w \subseteq \{-1, 0, 1\}$ or $\mathbf{dsfxs} w \subseteq \{0, 1, 2\}$, then $w \in X$.

Proof. Suppose $w \in \{0, 1\}^*$ and assume that either $\mathbf{dsfxs} w \subseteq \{-2, -1, 0\}$ or $\mathbf{dsfxs} w \subseteq \{-1, 0, 1\}$ or $\mathbf{dsfxs} w \subseteq \{0, 1, 2\}$. There are three case to consider.

- Suppose $\mathbf{dsfxs} w \subseteq \{-2, -1, 0\}$. Because $-2 \in \{-2, -1, 0\}$ and $\mathbf{dsfxs} w \subseteq \{-2, -1, 0\} = \{-2, (-2) + 1, (-2) + 2\}$, Lemma PS4.3.3 tells us that $w \in X$.
- Suppose $\mathbf{dsfxs} w \subseteq \{-1, 0, 1\}$. Because $-1 \in \{-2, -1, 0\}$ and $\mathbf{dsfxs} w \subseteq \{-1, 0, 1\} = \{-1, (-1) + 1, (-1) + 2\}$, Lemma PS4.3.3 tells us that $w \in X$.
- Suppose $\mathbf{dsfxs} w \subseteq \{0, 1, 2\}$. Because $0 \in \{-2, -1, 0\}$ and $\mathbf{dsfxs} w \subseteq \{0, 1, 2\} = \{0, 0+1, 0+2\}$, Lemma PS4.3.3 tells us that $w \in X$.

\square

For $-2 \leq n \leq 0 \leq m \leq 2$, define

$$Y^{n,m} = \{w \in \{0, 1\}^* \mid \mathbf{dsfxs} w \subseteq \{n, \dots, m\}\}.$$

Thus it is easy to show that:

- if v is a suffix of $w \in Y^{n,m}$, then $n \leq \mathbf{diff} v \leq m$;
- if $w \in \{0, 1\}^*$ and, for all suffixes v of w , $n \leq \mathbf{diff} v \leq m$, then $w \in Y^{n,m}$;
- $\% \in Y^{n,m}$.

The basis of the proof that $L(M) = X$ is the following lemma:

Lemma PS4.3.5

- (1) $X = Y^{0,2} \cup Y^{-1,1} \cup Y^{-2,0}$.
- (2) $Y^{0,2} = \{\%\} \cup Y^{-1,1}\{1\}$.
- (3) $Y^{-1,1} = \{\%\} \cup Y^{0,2}\{0\} \cup Y^{-2,0}\{1\}$.
- (4) $Y^{-2,0} = \{\%\} \cup Y^{-1,1}\{0\}$.

Proof.

- (1) We show that $X \subseteq Y^{0,2} \cup Y^{-1,1} \cup Y^{-2,0} \subseteq X$.

- To show $X \subseteq Y^{0,2} \cup Y^{-1,1} \cup Y^{-2,0}$, suppose $w \in X$. By Lemma PS4.3.2, we have that either $\mathbf{dsfxs} w \subseteq \{-2, -1, 0\}$ or $\mathbf{dsfxs} w \subseteq \{-1, 0, 1\}$ or $\mathbf{dsfxs} w \subseteq \{0, 1, 2\}$. Thus there are three cases to consider.

- Suppose $\mathbf{dsfxs} w \subseteq \{-2, -1, 0\}$. Thus $w \in Y^{-2,0} \subseteq Y^{0,2} \cup Y^{-1,1} \cup Y^{-2,0}$.
- Suppose $\mathbf{dsfxs} w \subseteq \{-1, 0, 1\}$. Thus $w \in Y^{-1,1} \subseteq Y^{0,2} \cup Y^{-1,1} \cup Y^{-2,0}$.
- Suppose $\mathbf{dsfxs} w \subseteq \{0, 1, 2\}$. Thus $w \in Y^{0,2} \subseteq Y^{0,2} \cup Y^{-1,1} \cup Y^{-2,0}$.
- To show $Y^{0,2} \cup Y^{-1,1} \cup Y^{-2,0} \subseteq X$, suppose $w \in Y^{0,2} \cup Y^{-1,1} \cup Y^{-2,0}$. There are three cases to consider.
 - Suppose $w \in Y^{0,2}$, so that $\mathbf{dsfxs} w \subseteq \{0, 1, 2\}$. Thus $w \in X$, by Lemma PS4.3.4.
 - Suppose $w \in Y^{-1,1}$, so that $\mathbf{dsfxs} w \subseteq \{-1, 0, 1\}$. Thus $w \in X$, by Lemma PS4.3.4.
 - Suppose $w \in Y^{-2,0}$, so that $\mathbf{dsfxs} w \subseteq \{-2, -1, 0\}$. Thus $w \in X$, by Lemma PS4.3.4.

(2) We show that $Y^{0,2} \subseteq \{\% \} \cup Y^{-1,1}\{1\} \subseteq Y^{0,2}$.

- To show that $Y^{0,2} \subseteq \{\% \} \cup Y^{-1,1}\{1\}$, suppose $w \in Y^{0,2}$. If $w = \%$, then $w \in \{\% \} \cup Y^{-1,1}\{1\}$. So, suppose $w \neq \%$. Then $w = xa$ for some $x \in \{0, 1\}^*$ and $a \in \{0, 1\}$. We cannot have $a = 0$, as then $-1 \in \mathbf{dsfxs} w$ (contradicting $w \in Y^{0,2}$). Thus $a = 1$, so that $w = x1$. To see that $x \in Y^{-1,1}$, suppose v is a suffix of x . Because $v1$ is a suffix of $w \in Y^{0,2}$, we have that $0 \leq \mathbf{diff}(v1) \leq 2$. But $\mathbf{diff}(v1) = \mathbf{diff} v + 1$, and thus $-1 \leq \mathbf{diff} v \leq 1$. Thus $w = x1 \in Y^{-1,1}\{1\} \subseteq \{\% \} \cup Y^{-1,1}\{1\}$.
- To show that $\{\% \} \cup Y^{-1,1}\{1\} \subseteq Y^{0,2}$, suppose $w \in \{\% \} \cup Y^{-1,1}\{1\}$. If $w \in \{\% \}$, then $w \in Y^{0,2}$. Otherwise, we have that $w \in Y^{-1,1}\{1\}$, so that $w = x1$, for some $x \in Y^{-1,1}$. To see that $w \in Y^{0,2}$, suppose v is a suffix of $w = x1$. We must show that $0 \leq \mathbf{diff} v \leq 2$. If $v = \%$, then this is true. Otherwise $v = u1$ for some suffix u of x . Because $x \in Y^{-1,1}$, we have that $-1 \leq \mathbf{diff} u \leq 1$. Thus $0 \leq \mathbf{diff} v \leq 2$.

(3) We show that $Y^{-1,1} \subseteq \{\% \} \cup Y^{0,2}\{0\} \cup Y^{-2,0}\{1\} \subseteq Y^{-1,1}$.

- To show that $Y^{-1,1} \subseteq \{\% \} \cup Y^{0,2}\{0\} \cup Y^{-2,0}\{1\}$, suppose $w \in Y^{-1,1}$. If $w = \%$, then $w \in \{\% \} \cup Y^{0,2}\{0\} \cup Y^{-2,0}\{1\}$. So, suppose $w \neq \%$. Then $w = xa$ for some $x \in \{0, 1\}^*$ and $a \in \{0, 1\}$. There are two cases to consider.
 - Suppose $a = 0$, so that $w = x0$. To see that $x \in Y^{0,2}$, suppose v is a suffix of x . Because $v0$ is a suffix of $w \in Y^{-1,1}$, we have that $-1 \leq \mathbf{diff}(v0) \leq 1$. But $\mathbf{diff}(v0) = \mathbf{diff} v + 1$, and thus $0 \leq \mathbf{diff} v \leq 2$. Thus $w = x0 \in Y^{0,2}\{0\} \subseteq \{\% \} \cup Y^{0,2}\{0\} \cup Y^{-2,0}\{1\}$.
 - Suppose $a = 1$, so that $w = x1$. To see that $x \in Y^{-2,0}$, suppose v is a suffix of x . Because $v1$ is a suffix of $w \in Y^{-1,1}$, we have that $-1 \leq \mathbf{diff}(v1) \leq 1$. But $\mathbf{diff}(v1) = \mathbf{diff} v + 1$, and thus $-2 \leq \mathbf{diff} v \leq 0$. Thus $w = x1 \in Y^{-2,0}\{1\} \subseteq \{\% \} \cup Y^{0,2}\{0\} \cup Y^{-2,0}\{1\}$.
- To show that $\{\% \} \cup Y^{0,2}\{0\} \cup Y^{-2,0}\{1\} \subseteq Y^{-1,1}$, suppose $w \in \{\% \} \cup Y^{0,2}\{0\} \cup Y^{-2,0}\{1\}$. If $w \in \{\% \}$, then $w \in Y^{-1,1}$. Otherwise, there are two cases to consider.
 - Suppose $w \in Y^{0,2}\{0\}$, so that $w = x0$, for some $x \in Y^{0,2}$. To see that $w \in Y^{-1,1}$, suppose v is a suffix of $w = x0$. We must show that $-1 \leq \mathbf{diff} v \leq 1$. If $v = \%$, then this is true. Otherwise $v = u0$ for some suffix u of x . Because $x \in Y^{0,2}$, we have that $0 \leq \mathbf{diff} u \leq 2$. Thus $-1 \leq \mathbf{diff} v \leq 1$.

- Suppose $w \in Y^{-2,0}\{1\}$, so that $w = x1$, for some $x \in Y^{-2,0}$. To see that $w \in Y^{-1,1}$, suppose v is a suffix of $w = x1$. We must show that $-1 \leq \mathbf{diff} v \leq 1$. If $v = \%$, then this is true. Otherwise $v = u1$ for some suffix u of x . Because $x \in Y^{-2,0}$, we have that $-2 \leq \mathbf{diff} u \leq 0$. Thus $-1 \leq \mathbf{diff} v \leq 1$.

(4) We show that $Y^{-2,0} \subseteq \{\%\} \cup Y^{-1,1}\{0\} \subseteq Y^{-2,0}$.

- To show that $Y^{-2,0} \subseteq \{\%\} \cup Y^{-1,1}\{0\}$, suppose $w \in Y^{-2,0}$. If $w = \%$, then $w \in \{\%\} \cup Y^{-1,1}\{0\}$. So, suppose $w \neq \%$. Then $w = xa$ for some $x \in \{0,1\}^*$ and $a \in \{0,1\}$. We cannot have $a = 1$, as then $1 \in \mathbf{dsfxs} w$ (contradicting $w \in Y^{-2,0}$). Thus $a = 0$, so that $w = x0$. To see that $x \in Y^{-1,1}$, suppose v is a suffix of x . Because $v0$ is a suffix of $w \in Y^{-2,0}$, we have that $-2 \leq \mathbf{diff}(v0) \leq 0$. But $\mathbf{diff}(v0) = \mathbf{diff} v + -1$, and thus $-1 \leq \mathbf{diff} v \leq 1$. Thus $w = x0 \in Y^{-1,1}\{0\} \subseteq \{\%\} \cup Y^{-1,1}\{0\}$.
- To show that $\{\%\} \cup Y^{-1,1}\{0\} \subseteq Y^{-2,0}$, suppose $w \in \{\%\} \cup Y^{-1,1}\{0\}$. If $w \in \{\%\}$, then $w \in Y^{-2,0}$. Otherwise, we have that $w \in Y^{-1,1}\{0\}$, so that $w = x0$, for some $x \in Y^{-1,1}$. To see that $w \in Y^{-2,0}$, suppose v is a suffix of $w = x0$. We must show that $-2 \leq \mathbf{diff} v \leq 0$. If $v = \%$, then this is true. Otherwise $v = u0$ for some suffix u of x . Because $x \in Y^{-1,1}$, we have that $-1 \leq \mathbf{diff} u \leq 1$. Thus $-2 \leq \mathbf{diff} v \leq 0$.

□

In what follows, we will show that $\Lambda_A = \{\%\}$, $\Lambda_B = Y^{0,2}$, $\Lambda_C = Y^{-1,1}$ and $\Lambda_D = Y^{-2,0}$.

Lemma PS4.3.6

- (A) For all $w \in \Lambda_A$, $w \in \{\%\}$.
- (B) For all $w \in \Lambda_B$, $w \in Y^{0,2}$.
- (C) For all $w \in \Lambda_C$, $w \in Y^{-1,1}$.
- (D) For all $w \in \Lambda_D$, $w \in Y^{-2,0}$.

Proof. We proceed by induction on Λ . There are 8 (1 plus the number of transitions) parts to show.

(empty string) Clearly $\% \in \{\%\}$, as required.

(A, % \rightarrow B) Suppose $w \in \Lambda_A$, and assume the inductive hypothesis: $w \in \{\%\}$. We must show that $w\% \in Y^{0,2}$. And $w\% = \%\% = \% \in Y^{0,2}$.

(A, % \rightarrow C) Suppose $w \in \Lambda_A$, and assume the inductive hypothesis: $w \in \{\%\}$. We must show that $w\% \in Y^{-1,1}$. And $w\% = \%\% = \% \in Y^{-1,1}$.

(A, % \rightarrow D) Suppose $w \in \Lambda_A$, and assume the inductive hypothesis: $w \in \{\%\}$. We must show that $w\% \in Y^{-2,0}$. And $w\% = \%\% = \% \in Y^{-2,0}$.

(B, 0 \rightarrow C) Suppose $w \in \Lambda_B$, and assume the inductive hypothesis: $w \in Y^{0,2}$. We must show that $w0 \in Y^{-1,1}$. And $w0 \in Y^{0,2}\{0\} \subseteq Y^{-1,1}$, by Lemma PS4.3.5(3).

(C, 0 \rightarrow D) Suppose $w \in \Lambda_C$, and assume the inductive hypothesis: $w \in Y^{-1,1}$. We must show that $w0 \in Y^{-2,0}$. And $w0 \in Y^{-1,1}\{0\} \subseteq Y^{-2,0}$, by Lemma PS4.3.5(4).

(C, 1 \rightarrow B) Suppose $w \in \Lambda_C$, and assume the inductive hypothesis: $w \in Y^{-1,1}$. We must show that $w1 \in Y^{0,2}$. And $w1 \in Y^{-1,1}\{1\} \subseteq Y^{0,2}$, by Lemma PS4.3.5(2).

(D, 1 \rightarrow C) Suppose $w \in \Lambda_D$, and assume the inductive hypothesis: $w \in Y^{-2,0}$. We must show that $w1 \in Y^{-1,1}$. And $w1 \in Y^{-2,0}\{1\} \subseteq Y^{-1,1}$, by Lemma PS4.3.5(3).

□

Lemma PS4.3.7

For all $w \in \{0, 1\}^*$:

- (A) if $w \in \{\%\}$, then $w \in \Lambda_A$;
- (B) if $w \in Y^{0,2}$, then $w \in \Lambda_B$;
- (C) if $w \in Y^{-1,1}$, then $w \in \Lambda_C$;
- (D) if $w \in Y^{-2,0}$, then $w \in \Lambda_D$.

Proof. We proceed by strong string induction. Suppose $w \in \{0, 1\}^*$, and assume the inductive hypothesis: for all $x \in \{0, 1\}^*$, if x is a proper substring of w , then

- (A) if $x \in \{\%\}$, then $x \in \Lambda_A$;
- (B) if $x \in Y^{0,2}$, then $x \in \Lambda_B$;
- (C) if $x \in Y^{-1,1}$, then $x \in \Lambda_C$;
- (D) if $x \in Y^{-2,0}$, then $x \in \Lambda_D$.

We must show that

- (A) if $w \in \{\%\}$, then $w \in \Lambda_A$;
- (B) if $w \in Y^{0,2}$, then $w \in \Lambda_B$;
- (C) if $w \in Y^{-1,1}$, then $w \in \Lambda_C$;
- (D) if $w \in Y^{-2,0}$, then $w \in \Lambda_D$.

There are four cases to consider.

- (A) Suppose $w \in \{\%\}$. We must show that $w \in \Lambda_A$. Because **A** is M 's start state, $w = \% \in \Lambda_A$.
- (B) Suppose $w \in Y^{0,2}$. We must show that $w \in \Lambda_B$. By Lemma PS4.3.5(2), we have that $w \in \{\%\} \cup Y^{-1,1}\{1\}$. Thus there are two subcases to consider.
 - Suppose $w \in \{\%\}$. Because **A** is M 's start state, we have $\% \in \Lambda_A$. And since $(A, \%, B) \in T_M$, it follows that $w = \% = \% \% \in \Lambda_B$.

- Suppose $w \in Y^{-1,1}\{1\}$, so that $w = x1$, for some $x \in Y^{-1,1}$. Because x is a proper substring of w , part (C) of the inductive hypothesis tells us that $x \in \Lambda_C$. Thus $w = x1 \in \Lambda_B$, because of the transition $(C, 1, B)$.
- (C) Suppose $w \in Y^{-1,1}$. We must show that $w \in \Lambda_C$. By Lemma PS4.3.5(3), we have that $w \in \{\% \} \cup Y^{0,2}\{0\} \cup Y^{-2,0}\{1\}$. Thus there are three subcases to consider.
- Suppose $w \in \{\% \}$. Because **A** is M 's start state, we have $\% \in \Lambda_A$. And since $(A, \%, C) \in T_M$, it follows that $w = \% = \% \% \in \Lambda_C$.
 - Suppose $w \in Y^{0,2}\{0\}$, so that $w = x0$, for some $x \in Y^{0,2}$. Because x is a proper substring of w , part (B) of the inductive hypothesis tells us that $x \in \Lambda_B$. Thus $w = x0 \in \Lambda_C$, because of the transition $(B, 0, C)$.
 - Suppose $w \in Y^{-2,0}\{1\}$, so that $w = x1$, for some $x \in Y^{-2,0}$. Because x is a proper substring of w , part (D) of the inductive hypothesis tells us that $x \in \Lambda_D$. Thus $w = x1 \in \Lambda_C$, because of the transition $(D, 1, C)$.
- (D) Suppose $w \in Y^{-2,0}$. We must show that $w \in \Lambda_D$. By Lemma PS4.3.5(4), we have that $w \in \{\% \} \cup Y^{-1,1}\{0\}$. Thus there are two subcases to consider.
- Suppose $w \in \{\% \}$. Because **A** is M 's start state, we have $\% \in \Lambda_A$. And since $(A, \%, D) \in T_M$, it follows that $w = \% = \% \% \in \Lambda_D$.
 - Suppose $w \in Y^{-1,1}\{0\}$, so that $w = x0$, for some $x \in Y^{-1,1}$. Because x is a proper substring of w , part (C) of the inductive hypothesis tells us that $x \in \Lambda_C$. Thus $w = x0 \in \Lambda_D$, because of the transition $(C, 0, D)$.

□

Lemma PS4.3.8

- (A) $\Lambda_A = \{\% \}$.
- (B) $\Lambda_B = Y^{0,2}$.
- (C) $\Lambda_C = Y^{-1,1}$.
- (D) $\Lambda_D = Y^{-2,0}$.

Proof. Follows by Lemmas PS4.3.6 and PS4.3.7. □

Lemma PS4.3.9

$L(M) = X$.

Proof. Because M 's set of accepting states is $\{B, C, D\}$, it follows that $L(M) = \Lambda_B \cup \Lambda_C \cup \Lambda_D$. And by Lemma PS4.3.8 and Lemma PS4.3.5(1), we have that $\Lambda_B \cup \Lambda_C \cup \Lambda_D = Y^{0,2} \cup Y^{-1,1} \cup Y^{-2,0} = X$. Thus $L(M) = X$. □