

## *4.5: Proving the Correctness of Grammars*

In this section, we consider techniques for proving the correctness of grammars, i.e., for proving that grammars generate the languages we want them to.

## *Definition of $\Pi$*

Suppose  $G$  is a grammar and  $a \in Q_G \cup \mathbf{alphabet} G$ . Then  $\Pi_{G,a} = \{ w \in (\mathbf{alphabet} G)^* \mid w \text{ is parsable from } a \text{ using } G \}$ .

If it's clear which grammar we are talking about, we often abbreviate  $\Pi_{G,a}$  to  $\Pi_a$ .

Clearly,  $\Pi_{G,s_G} =$

## *Definition of $\Pi$*

Suppose  $G$  is a grammar and  $a \in Q_G \cup \mathbf{alphabet} G$ . Then  $\Pi_{G,a} = \{ w \in (\mathbf{alphabet} G)^* \mid w \text{ is parsable from } a \text{ using } G \}$ .

If it's clear which grammar we are talking about, we often abbreviate  $\Pi_{G,a}$  to  $\Pi_a$ .

Clearly,  $\Pi_{G,s_G} = L(G)$ .

## Definition of $\Pi$

Suppose  $G$  is a grammar and  $a \in Q_G \cup \mathbf{alphabet} G$ . Then  $\Pi_{G,a} = \{ w \in (\mathbf{alphabet} G)^* \mid w \text{ is parsable from } a \text{ using } G \}$ .

If it's clear which grammar we are talking about, we often abbreviate  $\Pi_{G,a}$  to  $\Pi_a$ .

Clearly,  $\Pi_{G,s_G} = L(G)$ .

For example, if  $G$  is the grammar

$$A \rightarrow \% \mid 0A1$$

then  $\Pi_0 =$

## Definition of $\Pi$

Suppose  $G$  is a grammar and  $a \in Q_G \cup \mathbf{alphabet} G$ . Then  $\Pi_{G,a} = \{ w \in (\mathbf{alphabet} G)^* \mid w \text{ is parsable from } a \text{ using } G \}$ .

If it's clear which grammar we are talking about, we often abbreviate  $\Pi_{G,a}$  to  $\Pi_a$ .

Clearly,  $\Pi_{G,s_G} = L(G)$ .

For example, if  $G$  is the grammar

$$A \rightarrow \% \mid 0A1$$

then  $\Pi_0 = \{0\}$

## Definition of $\Pi$

Suppose  $G$  is a grammar and  $a \in Q_G \cup \mathbf{alphabet} G$ . Then  $\Pi_{G,a} = \{ w \in (\mathbf{alphabet} G)^* \mid w \text{ is parsable from } a \text{ using } G \}$ .

If it's clear which grammar we are talking about, we often abbreviate  $\Pi_{G,a}$  to  $\Pi_a$ .

Clearly,  $\Pi_{G,s_G} = L(G)$ .

For example, if  $G$  is the grammar

$$A \rightarrow \% \mid 0A1$$

then  $\Pi_0 = \{0\}$ ,  $\Pi_1 = \{1\}$  and  $\Pi_A =$

## Definition of $\Pi$

Suppose  $G$  is a grammar and  $a \in Q_G \cup \mathbf{alphabet} G$ . Then  $\Pi_{G,a} = \{ w \in (\mathbf{alphabet} G)^* \mid w \text{ is parsable from } a \text{ using } G \}$ .

If it's clear which grammar we are talking about, we often abbreviate  $\Pi_{G,a}$  to  $\Pi_a$ .

Clearly,  $\Pi_{G,s_G} = L(G)$ .

For example, if  $G$  is the grammar

$$A \rightarrow \% \mid 0A1$$

then  $\Pi_0 = \{0\}$ ,  $\Pi_1 = \{1\}$  and  $\Pi_A = \{0^n 1^n \mid n \in \mathbb{N}\} =$

## Definition of $\Pi$

Suppose  $G$  is a grammar and  $a \in Q_G \cup \text{alphabet } G$ . Then  $\Pi_{G,a} = \{ w \in (\text{alphabet } G)^* \mid w \text{ is parsable from } a \text{ using } G \}$ .

If it's clear which grammar we are talking about, we often abbreviate  $\Pi_{G,a}$  to  $\Pi_a$ .

Clearly,  $\Pi_{G,s_G} = L(G)$ .

For example, if  $G$  is the grammar

$$A \rightarrow \% \mid 0A1$$

then  $\Pi_0 = \{0\}$ ,  $\Pi_1 = \{1\}$  and  $\Pi_A = \{0^n 1^n \mid n \in \mathbb{N}\} = L(G)$ .



## Properties of $\Pi$

### Proposition 4.5.1

Suppose  $G$  is a grammar.

- (1) For all  $a \in \mathbf{alphabet } G$ ,  $\Pi_{G,a} =$
- (2) For all  $q \in Q_G$ , if  $q \rightarrow \% \in P_G$ , then  $\Pi_{G,q} =$
- (3) For all  $q \in Q_G$ ,  $n \in \mathbb{N} - \{0\}$ ,  $a_1, \dots, a_n \in \mathbf{Sym}$  and  $w_1, \dots, w_n \in \mathbf{Str}$ , if  $q \rightarrow a_1 \cdots a_n \in P_G$  and  $w_1 \in \Pi_{G,a_1}, \dots, w_n \in \Pi_{G,a_n}$ , then  $\Pi_{G,q} =$

## Properties of $\Pi$

### Proposition 4.5.1

Suppose  $G$  is a grammar.

- (1) For all  $a \in \mathbf{alphabet } G$ ,  $\Pi_{G,a} = \{a\}$ .
- (2) For all  $q \in Q_G$ , if  $q \rightarrow \% \in P_G$ , then  $\% \in \Pi_{G,q}$ .
- (3) For all  $q \in Q_G$ ,  $n \in \mathbb{N} - \{0\}$ ,  $a_1, \dots, a_n \in \mathbf{Sym}$  and  $w_1, \dots, w_n \in \mathbf{Str}$ , if  $q \rightarrow a_1 \cdots a_n \in P_G$  and  $w_1 \in \Pi_{G,a_1}, \dots, w_n \in \Pi_{G,a_n}$ , then  $w_1 \cdots w_n \in \Pi_{G,q}$ .

## Properties of $\Pi$

### Proposition 4.5.1

Suppose  $G$  is a grammar.

- (1) For all  $a \in \mathbf{alphabet } G$ ,  $\Pi_{G,a} = \{a\}$ .
- (2) For all  $q \in Q_G$ , if  $q \rightarrow \% \in P_G$ , then  $\% \in \Pi_{G,q}$ .
- (3) For all  $q \in Q_G$ ,  $n \in \mathbb{N} - \{0\}$ ,  $a_1, \dots, a_n \in \mathbf{Sym}$  and  $w_1, \dots, w_n \in \mathbf{Str}$ , if  $q \rightarrow a_1 \cdots a_n \in P_G$  and  $w_1 \in \Pi_{G,a_1}, \dots, w_n \in \Pi_{G,a_n}$ , then  $w_1 \cdots w_n \in \Pi_{G,q}$ .

## Properties of $\Pi$

### Proposition 4.5.1

Suppose  $G$  is a grammar.

- (1) For all  $a \in \mathbf{alphabet } G$ ,  $\Pi_{G,a} = \{a\}$ .
- (2) For all  $q \in Q_G$ , if  $q \rightarrow \% \in P_G$ , then  $\% \in \Pi_{G,q}$ .
- (3) For all  $q \in Q_G$ ,  $n \in \mathbb{N} - \{0\}$ ,  $a_1, \dots, a_n \in \mathbf{Sym}$  and  $w_1, \dots, w_n \in \mathbf{Str}$ , if  $q \rightarrow a_1 \cdots a_n \in P_G$  and  $w_1 \in \Pi_{G,a_1}, \dots, w_n \in \Pi_{G,a_n}$ , then  $w_1 \cdots w_n \in \Pi_{G,q}$ .

## Main Example

Define  $\mathbf{diff} \in \{0,1\}^* \rightarrow \mathbb{Z}$  by: for all  $w \in \{0,1\}^*$ ,

$\mathbf{diff} w =$  the number of 1's in  $w$   $-$  the number of 0's in  $w$ .

Then:

- $\mathbf{diff} \% = 0$ ;
- $\mathbf{diff} 1 = 1$ ;
- $\mathbf{diff} 0 = -1$ ; and
- for all  $x, y \in \{0,1\}^*$ ,  $\mathbf{diff}(xy) = \mathbf{diff} x + \mathbf{diff} y$ .

## *Main Example*

Our main example will be the grammar  $G$ :

$$A \rightarrow \% \mid 0BA \mid 1CA,$$

$$B \rightarrow 1 \mid 0BB,$$

$$C \rightarrow 0 \mid 1CC.$$

Let

$$X = \{ w \in \{0,1\}^* \mid \mathbf{diff} w = 0 \},$$

$$Y = \{ w \in \{0,1\}^* \mid \mathbf{diff} w = 1 \text{ and,}$$

$$Z = \{ w \in \{0,1\}^* \mid \mathbf{diff} w = -1 \text{ and,}$$

We will prove that  $L(G) = \Pi_{G,A} = X$ ,  $\Pi_{G,B} = Y$  and  $\Pi_{G,C} = Z$ .

## Main Example

Our main example will be the grammar  $G$ :

$$A \rightarrow 0 \mid 0BA \mid 1CA,$$

$$B \rightarrow 1 \mid 0BB,$$

$$C \rightarrow 0 \mid 1CC.$$

Let

$$X = \{ w \in \{0,1\}^* \mid \mathbf{diff} w = 0 \},$$

$$Y = \{ w \in \{0,1\}^* \mid \mathbf{diff} w = 1 \text{ and,} \\ \text{for all proper prefixes } v \text{ of } w, \mathbf{diff} v \leq 0 \},$$

$$Z = \{ w \in \{0,1\}^* \mid \mathbf{diff} w = -1 \text{ and,} \\ \text{for all proper prefixes } v \text{ of } w, \mathbf{diff} v \geq 0 \}.$$

We will prove that  $L(G) = \Pi_{G,A} = X$ ,  $\Pi_{G,B} = Y$  and  $\Pi_{G,C} = Z$ .

## Main Example

### Lemma 4.5.2

Suppose  $x \in \{0, 1\}^*$ .

- (1) If  $\mathbf{diff} x \geq 1$ , then  $x = yz$  for some  $y, z \in \{0, 1\}^*$  such that  $y \in Y$  and  $\mathbf{diff} z = \mathbf{diff} x - 1$ .
- (2) If  $\mathbf{diff} x \leq -1$ , then  $x = yz$  for some  $y, z \in \{0, 1\}^*$  such that  $y \in Z$  and  $\mathbf{diff} z = \mathbf{diff} x + 1$ .

The proof is in the book.



## *Proving that Enough is Generated*

First we study techniques for showing that everything we want a grammar to generate is really generated.

Since  $X, Y, Z \subseteq \{0, 1\}^*$ , to prove that  $X \subseteq \Pi_{G,A}$ ,  $Y \subseteq \Pi_{G,B}$  and  $Z \subseteq \Pi_{G,C}$ , it will suffice to use strong string induction to show that, for all  $w \in \{0, 1\}^*$ :

- (A) if  $w \in X$ , then  $w \in \Pi_{G,A}$ ;
- (B) if  $w \in Y$ , then  $w \in \Pi_{G,B}$ ; and
- (C) if  $w \in Z$ , then  $w \in \Pi_{G,C}$ .

## *Enough is Generated in Example*

We proceed by strong string induction. Suppose  $w \in \{0, 1\}^*$ , and assume the inductive hypothesis: for all  $x \in \{0, 1\}^*$ , if  $x$  is a proper substring of  $w$ , then:

- (A) if  $x \in X$ , then  $x \in \Pi_A$ ;
- (B) if  $x \in Y$ , then  $x \in \Pi_B$ ; and
- (C) if  $x \in Z$ , then  $x \in \Pi_C$ .

We must prove that:

- (A) if  $w \in X$ , then  $w \in \Pi_A$ ;
- (B) if  $w \in Y$ , then  $w \in \Pi_B$ ; and
- (C) if  $w \in Z$ , then  $w \in \Pi_C$ .

## *Enough is Generated in Example*

(A) Suppose  $w \in X$ . We must show that  $w \in \Pi_A$ . There are three cases to consider.

- Suppose  $w = \%$ . Because  $A \rightarrow \% \in P$ , we have that  $w = \% \in \Pi_A$ .

## *Enough is Generated in Example*

(A) Suppose  $w \in X$ . We must show that  $w \in \Pi_A$ . There are three cases to consider.

- Suppose  $w = 0$ . Because  $A \rightarrow 0 \in P$ , we have that  $w = 0 \in \Pi_A$ .
- Suppose  $w = 0x$ , for some  $x \in \{0, 1\}^*$ . Because  $-1 + \mathbf{diff} x = \mathbf{diff} w = 0$ , we have that  $\mathbf{diff} x = 1$ . Thus, by Lemma 4.5.2(1), we have that  $x = yz$ , for some  $y, z \in \{0, 1\}^*$  such that  $y \in Y$  and  $\mathbf{diff} z = \mathbf{diff} x - 1 = 1 - 1 = 0$ . Thus  $w = 0yz$ ,  $y \in Y$  and  $z \in X$ . We have  $0 \in \Pi_0$ . Because  $y \in Y$  and  $z \in X$

## *Enough is Generated in Example*

(A) Suppose  $w \in X$ . We must show that  $w \in \Pi_A$ . There are three cases to consider.

- Suppose  $w = \%$ . Because  $A \rightarrow \% \in P$ , we have that  $w = \% \in \Pi_A$ .
- Suppose  $w = 0x$ , for some  $x \in \{0, 1\}^*$ . Because  $-1 + \mathbf{diff} x = \mathbf{diff} w = 0$ , we have that  $\mathbf{diff} x = 1$ . Thus, by Lemma 4.5.2(1), we have that  $x = yz$ , for some  $y, z \in \{0, 1\}^*$  such that  $y \in Y$  and  $\mathbf{diff} z = \mathbf{diff} x - 1 = 1 - 1 = 0$ . Thus  $w = 0yz$ ,  $y \in Y$  and  $z \in X$ . We have  $0 \in \Pi_0$ . Because  $y \in Y$  and  $z \in X$  are proper substrings of  $w$ , parts (B) and (A) of the inductive hypothesis tell us that  $y \in \Pi_B$  and  $z \in \Pi_A$ . Thus, because  $A \rightarrow 0BA \in P$ , it follows that that  $w = 0yz \in \Pi_A$ .

## *Enough is Generated in Example*

(A) Suppose  $w \in X$ . We must show that  $w \in \Pi_A$ . There are three cases to consider.

- Suppose  $w = \%$ . Because  $A \rightarrow \% \in P$ , we have that  $w = \% \in \Pi_A$ .
- Suppose  $w = 0x$ , for some  $x \in \{0, 1\}^*$ . Because  $-1 + \mathbf{diff} x = \mathbf{diff} w = 0$ , we have that  $\mathbf{diff} x = 1$ . Thus, by Lemma 4.5.2(1), we have that  $x = yz$ , for some  $y, z \in \{0, 1\}^*$  such that  $y \in Y$  and  $\mathbf{diff} z = \mathbf{diff} x - 1 = 1 - 1 = 0$ . Thus  $w = 0yz$ ,  $y \in Y$  and  $z \in X$ . We have  $0 \in \Pi_0$ . Because  $y \in Y$  and  $z \in X$  are proper substrings of  $w$ , parts (B) and (A) of the inductive hypothesis tell us that  $y \in \Pi_B$  and  $z \in \Pi_A$ . Thus, because  $A \rightarrow 0BA \in P$ , it follows that that  $w = 0yz \in \Pi_A$ .
- Suppose  $w = 1x$ , for some  $x \in \{0, 1\}^*$ . The proof is analogous to the preceding case.

## *Enough is Generated in Example*

(B) Suppose  $w \in Y$ . We must show that  $w \in \Pi_B$ . Because **diff**  $w = 1$ , there are two cases to consider.

- Suppose  $w = 1x$ , for some  $x \in \{0, 1\}^*$ .

## *Enough is Generated in Example*

- (B) Suppose  $w \in Y$ . We must show that  $w \in \Pi_B$ . Because  $\text{diff } w = 1$ , there are two cases to consider.
- Suppose  $w = 1x$ , for some  $x \in \{0, 1\}^*$ . Because all proper prefixes of  $w$  have  $\text{diffs} \leq 0$ , we have that



## *Enough is Generated in Example*

- (B) Suppose  $w \in Y$ . We must show that  $w \in \Pi_B$ . Because  $\text{diff } w = 1$ , there are two cases to consider.
- Suppose  $w = 1x$ , for some  $x \in \{0, 1\}^*$ . Because all proper prefixes of  $w$  have  $\text{diffs} \leq 0$ , we have that  $x = \epsilon$ , so that  $w = 1$ . Since  $B \rightarrow 1 \in P$ , we have that  $w = 1 \in \Pi_B$ .

## *Enough is Generated in Example*

(B) Suppose  $w \in Y$ . We must show that  $w \in \Pi_B$ . Because  $\text{diff } w = 1$ , there are two cases to consider.

- Suppose  $w = 1x$ , for some  $x \in \{0, 1\}^*$ . Because all proper prefixes of  $w$  have diffs  $\leq 0$ , we have that  $x = \%$ , so that  $w = 1$ . Since  $B \rightarrow 1 \in P$ , we have that  $w = 1 \in \Pi_B$ .
- Suppose  $w = 0x$ , for some  $x \in \{0, 1\}^*$ . Thus  $\text{diff } x = 2$ . Because  $\text{diff } x \geq 1$ , by Lemma 4.5.2(1), we have that  $x = yz$ , for some  $y, z \in \{0, 1\}^*$  such that  $y \in Y$  and  $\text{diff } z = \text{diff } x - 1 = 2 - 1 = 1$ . Hence  $w = 0yz$ .

## *Enough is Generated in Example*

(B) Suppose  $w \in Y$ . We must show that  $w \in \Pi_B$ . Because  $\text{diff } w = 1$ , there are two cases to consider.

- Suppose  $w = 1x$ , for some  $x \in \{0, 1\}^*$ . Because all proper prefixes of  $w$  have diffs  $\leq 0$ , we have that  $x = \%$ , so that  $w = 1$ . Since  $B \rightarrow 1 \in P$ , we have that  $w = 1 \in \Pi_B$ .
- Suppose  $w = 0x$ , for some  $x \in \{0, 1\}^*$ . Thus  $\text{diff } x = 2$ . Because  $\text{diff } x \geq 1$ , by Lemma 4.5.2(1), we have that  $x = yz$ , for some  $y, z \in \{0, 1\}^*$  such that  $y \in Y$  and  $\text{diff } z = \text{diff } x - 1 = 2 - 1 = 1$ . Hence  $w = 0yz$ . To finish the proof that  $z \in Y$ , suppose  $v$  is a proper prefix of  $z$ . Thus  $0yv$  is a proper prefix of  $w$ . Since  $w \in Y$ , it follows that  $\text{diff } v = \text{diff}(0yv) \leq 0$ , as required. Since  $y, z \in Y$ , part (B) of the inductive hypothesis tell us that  $y, z \in \Pi_B$ . Thus, because  $B \rightarrow 0BB \in P$  we have that  $w = 0yz \in \Pi_B$ .

## *Enough is Generated in Example*

(B) Suppose  $w \in Y$ . We must show that  $w \in \Pi_B$ . Because  $\text{diff } w = 1$ , there are two cases to consider.

- Suppose  $w = 1x$ , for some  $x \in \{0, 1\}^*$ . Because all proper prefixes of  $w$  have diffs  $\leq 0$ , we have that  $x = \epsilon$ , so that  $w = 1$ . Since  $B \rightarrow 1 \in P$ , we have that  $w = 1 \in \Pi_B$ .
- Suppose  $w = 0x$ , for some  $x \in \{0, 1\}^*$ . Thus  $\text{diff } x = 2$ . Because  $\text{diff } x \geq 1$ , by Lemma 4.5.2(1), we have that  $x = yz$ , for some  $y, z \in \{0, 1\}^*$  such that  $y \in Y$  and  $\text{diff } z = \text{diff } x - 1 = 2 - 1 = 1$ . Hence  $w = 0yz$ . To finish the proof that  $z \in Y$ , suppose  $v$  is a proper prefix of  $z$ . Thus  $0yv$  is a proper prefix of  $w$ . Since  $w \in Y$ , it follows that  $\text{diff } v = \text{diff}(0yv) \leq 0$ , as required. Since  $y, z \in Y$ , part (B) of the inductive hypothesis tell us that  $y, z \in \Pi_B$ . Thus, because  $B \rightarrow 0BB \in P$  we have that  $w = 0yz \in \Pi_B$ .

(C) Suppose  $w \in Z$ . We must show that  $w \in \Pi_C$ . The proof is analogous to the proof of part (B).

## *A Problem with Unit Productions*

Suppose  $H$  is the grammar

$$A \rightarrow B \mid 0A3, \quad B \rightarrow \% \mid 1B2,$$

and let

$$X = \{0^n 1^m 2^m 3^n \mid n, m \in \mathbb{N}\} \quad \text{and} \quad Y = \{1^m 2^m \mid m \in \mathbb{N}\}.$$

## *A Problem with Unit Productions*

Suppose  $H$  is the grammar

$$A \rightarrow B \mid 0A3, \quad B \rightarrow \% \mid 1B2,$$

and let

$$X = \{0^n 1^m 2^m 3^n \mid n, m \in \mathbb{N}\} \quad \text{and} \quad Y = \{1^m 2^m \mid m \in \mathbb{N}\}.$$

We can prove that  $X \subseteq \Pi_{H,A} = L(H)$  and  $Y \subseteq \Pi_{H,B}$  using the above technique, but the production  $A \rightarrow B$ , which is called a *unit production* because its right side is a single variable, makes part (A) tricky. If  $w = 0^0 1^m 2^m 3^0 = 1^m 2^m \in Y$ , we would like to use part (B) of the inductive hypothesis to conclude  $w \in \Pi_B$ , and then use the fact that  $A \rightarrow B \in P$  to conclude that  $w \in \Pi_A$ .

## A Problem with Unit Productions

Suppose  $H$  is the grammar

$$A \rightarrow B \mid 0A3, \quad B \rightarrow \% \mid 1B2,$$

and let

$$X = \{0^n 1^m 2^m 3^n \mid n, m \in \mathbb{N}\} \quad \text{and} \quad Y = \{1^m 2^m \mid m \in \mathbb{N}\}.$$

We can prove that  $X \subseteq \Pi_{H,A} = L(H)$  and  $Y \subseteq \Pi_{H,B}$  using the above technique, but the production  $A \rightarrow B$ , which is called a *unit production* because its right side is a single variable, makes part (A) tricky. If  $w = 0^0 1^m 2^m 3^0 = 1^m 2^m \in Y$ , we would like to use part (B) of the inductive hypothesis to conclude  $w \in \Pi_B$ , and then use the fact that  $A \rightarrow B \in P$  to conclude that  $w \in \Pi_A$ . But  $w$  is not a proper substring of itself, and so the inductive hypothesis is not applicable. Instead, we must split into cases  $m = 0$  and  $m \geq 1$ , using  $A \rightarrow B$  and  $B \rightarrow \%$ , in the first case, and  $A \rightarrow B$  and  $B \rightarrow 1B2$ , as well as the inductive hypothesis on  $1^{m-1} 2^{m-1} \in Y$ , in the second case.

## *A Problem with Unit Productions*

Because there are no productions from **B** back to **A**, we could also first use strong string induction to prove that, for all  $w \in \{0, 1\}^*$ ,

(B) if  $w \in Y$ , then  $w \in \Pi_B$ ,

and then use the result of this induction along with strong string induction to prove that for all  $w \in \{0, 1\}^*$ ,

(A) if  $w \in X$ , then  $w \in \Pi_A$ .

This works whenever two parts of a grammar are not mutually recursive.



## *A Problem with Unit Productions*

Because there are no productions from **B** back to **A**, we could also first use strong string induction to prove that, for all  $w \in \{0, 1\}^*$ ,

(B) if  $w \in Y$ , then  $w \in \Pi_B$ ,

and then use the result of this induction along with strong string induction to prove that for all  $w \in \{0, 1\}^*$ ,

(A) if  $w \in X$ , then  $w \in \Pi_A$ .

This works whenever two parts of a grammar are not mutually recursive.

With this grammar, we could also first use mathematical induction to prove that, for all  $m \in \mathbb{N}$ ,  $1^m 2^m \in \Pi_B$ , and then use the result of this induction to prove, by mathematical induction on  $n$ , that for all  $n, m \in \mathbb{N}$ ,  $0^n 1^m 2^m 3^n \in \Pi_A$ .

## *A Problem with $\epsilon$ -Productions*

Note that  $\epsilon$ -productions, i.e., productions of the form  $\alpha \rightarrow \epsilon$ , can cause similar problems to those caused by unit productions. E.g., if we have the productions

$$A \rightarrow BC \quad \text{and} \quad B \rightarrow \epsilon,$$

then  $A \rightarrow BC$  behaves like a unit production.

## *Proving that Everything Generated is Wanted*

To prove that everything generated by a grammar is wanted, we introduce a new induction principle that we call induction on  $\Pi$ .

## Principle of Induction on $\Pi$

### Theorem 4.5.3 (Principle of Induction on $\Pi$ )

Suppose  $G$  is a grammar,  $P_q(w)$  is a property of a string  $w \in \Pi_{G,q}$ , for all  $q \in Q_G$ , and  $P_a(w)$ , for  $a \in \mathbf{alphabet\ } G$ , says “ $w = a$ ”.

If

- (1) for all  $q \in Q_G$ , if  $q \rightarrow \% \in P_G$ , then  $P_q(\%)$ , and
- (2) for all  $q \in Q_G$ ,  $n \in \mathbb{N} - \{0\}$ ,  $a_1, \dots, a_n \in Q_G \cup \mathbf{alphabet\ } G$ , and  $w_1 \in \Pi_{G,a_1}, \dots, w_n \in \Pi_{G,a_n}$ , if  $q \rightarrow a_1 \cdots a_n \in P_G$  and  
( $\dagger$ ) then  $P_q(w_1 \cdots w_n)$ ,

then

for all  $q \in Q_G$ , for all  $w \in \Pi_{G,q}$ ,  $P_q(w)$ .

We refer to ( $\dagger$ ) as the inductive hypothesis.

## Principle of Induction on $\Pi$

### Theorem 4.5.3 (Principle of Induction on $\Pi$ )

Suppose  $G$  is a grammar,  $P_q(w)$  is a property of a string  $w \in \Pi_{G,q}$ , for all  $q \in Q_G$ , and  $P_a(w)$ , for  $a \in \mathbf{alphabet\ } G$ , says “ $w = a$ ”.

If

- (1) for all  $q \in Q_G$ , if  $q \rightarrow \% \in P_G$ , then  $P_q(\%)$ , and
- (2) for all  $q \in Q_G$ ,  $n \in \mathbb{N} - \{0\}$ ,  $a_1, \dots, a_n \in Q_G \cup \mathbf{alphabet\ } G$ , and  $w_1 \in \Pi_{G,a_1}, \dots, w_n \in \Pi_{G,a_n}$ , if  $q \rightarrow a_1 \cdots a_n \in P_G$  and  
(†)  $P_{a_1}(w_1), \dots, P_{a_n}(w_n)$ , then  $P_q(w_1 \cdots w_n)$ ,

then

for all  $q \in Q_G$ , for all  $w \in \Pi_{G,q}$ ,  $P_q(w)$ .

We refer to (†) as the inductive hypothesis.

## *Principle of Induction on $\Pi$*

**Proof.** It suffices to show that, for all  $pt \in \mathbf{PT}$ , for all  $q \in Q_G$  and  $w \in (\mathbf{alphabet } G)^*$ , if  $pt$  is valid for  $G$ ,  $\mathbf{rootLabel } pt = q$  and  $\mathbf{yield } pt = w$ , then  $P_q(w)$ . We prove this using the principle of induction on parse trees.  $\square$

## *Principle of Induction on $\Pi$*

**Proof.** It suffices to show that, for all  $pt \in \mathbf{PT}$ , for all  $q \in Q_G$  and  $w \in (\mathbf{alphabet } G)^*$ , if  $pt$  is valid for  $G$ ,  $\mathbf{rootLabel } pt = q$  and  $\mathbf{yield } pt = w$ , then  $P_q(w)$ . We prove this using the principle of induction on parse trees.  $\square$

When proving part (2), we can make use of the fact that, for  $a_i \in \mathbf{alphabet } G$ ,  $\Pi_{a_i} = \{a_i\}$ , so that  $w_i \in \Pi_{a_i}$  will be  $a_i$ . Hence it will be unnecessary to assume that  $P_{a_i}(a_i)$ , since this says " $a_i = a_i$ ", and so is always true.

## Using Induction on $\Pi$ in Example

Consider, again, our main example grammar  $G$ :

$$A \rightarrow \epsilon \mid 0BA \mid 1CA,$$

$$B \rightarrow 1 \mid 0BB,$$

$$C \rightarrow 0 \mid 1CC.$$

Let

$$X = \{ w \in \{0,1\}^* \mid \mathbf{diff} w = 0 \},$$

$$Y = \{ w \in \{0,1\}^* \mid \mathbf{diff} w = 1 \text{ and,}$$

for all proper prefixes  $v$  of  $w$ ,  $\mathbf{diff} v \leq 0 \}$ , and

$$Z = \{ w \in \{0,1\}^* \mid \mathbf{diff} w = -1 \text{ and,}$$

for all proper prefixes  $v$  of  $w$ ,  $\mathbf{diff} v \geq 0 \}$ .



## Using Induction on $\Pi$ in Example

We have already proven that  $X \subseteq \Pi_A = L(G)$ ,  $Y \subseteq \Pi_B$  and  $Z \subseteq \Pi_C$ . To complete the proof that  $L(G) = \Pi_A = X$ ,  $\Pi_B = Y$  and  $\Pi_C = Z$ , we will use induction on  $\Pi$  to prove that  $\Pi_A \subseteq X$ ,  $\Pi_B \subseteq Y$  and  $\Pi_C \subseteq Z$ .

We use induction on  $\Pi$  to show that:

- (A) for all  $w \in \Pi_A$ ,  $w \in X$ ;
- (B) for all  $w \in \Pi_B$ ,  $w \in Y$ ; and
- (C) for all  $w \in \Pi_C$ ,  $w \in Z$ .

Formally, this means that we let the properties  $P_A(w)$ ,  $P_B(w)$  and  $P_C(w)$  be “ $w \in X$ ”, “ $w \in Y$ ” and “ $w \in Z$ ”, respectively, and then use the induction principle to prove that, for all  $q \in Q_G$ , for all  $w \in \Pi_q$ ,  $P_q(w)$ . But we will actually work more informally.

## *Using Induction on $\Pi$ in Example*

There are seven productions to consider.

- $(A \rightarrow \%)$  We must show that  $\% \in X$  (as “ $w \in X$ ” is the property of part (A)). And this holds since **diff**  $\% = 0$ .

## Using Induction on $\Pi$ in Example

There are seven productions to consider.

- $(A \rightarrow \%)$  We must show that  $\% \in X$  (as “ $w \in X$ ” is the property of part (A)). And this holds since  $\text{diff } \% = 0$ .
- $(A \rightarrow 0BA)$  Suppose  $w_1 \in \Pi_B$  and  $w_2 \in \Pi_A$  (as  $0BA$  is the right-side of the production, and  $0$  is in  $G$ 's alphabet), and assume the inductive hypothesis,  $w_1 \in Y$  (as this is the property of part (B)) and  $w_2 \in X$  (as this is the property of part (A)). We must show that  $0w_1w_2 \in X$ , as the production shows that  $0w_1w_2 \in \Pi_A$ . Because  $w_1 \in Y$  and  $w_2 \in X$ , we have that  $\text{diff } w_1 = 1$  and  $\text{diff } w_2 = 0$ . Thus  $\text{diff}(0w_1w_2) = -1 + 1 + 0 = 0$ , showing that  $0w_1w_2 \in X$ .

## Using Induction on $\Pi$ in Example

- **(B  $\rightarrow$  0BB)** Suppose  $w_1, w_2 \in \Pi_B$ , and assume the inductive hypothesis,  $w_1, w_2 \in Y$ . Thus  $w_1$  and  $w_2$  are nonempty. We must show that  $0w_1w_2 \in Y$ . Clearly,  $\mathbf{diff}(0w_1w_2) = -1 + 1 + 1 = 1$ . So, suppose  $v$  is a proper prefix of  $0w_1w_2$ . We must show that  $\mathbf{diff} v \leq 0$ .

## Using Induction on $\Pi$ in Example

- **(B  $\rightarrow$  0BB)** Suppose  $w_1, w_2 \in \Pi_B$ , and assume the inductive hypothesis,  $w_1, w_2 \in Y$ . Thus  $w_1$  and  $w_2$  are nonempty. We must show that  $0w_1w_2 \in Y$ . Clearly,  $\text{diff}(0w_1w_2) = -1 + 1 + 1 = 1$ . So, suppose  $v$  is a proper prefix of  $0w_1w_2$ . We must show that  $\text{diff } v \leq 0$ . There are three cases to consider.
  - Suppose  $v = \%$ . Then  $\text{diff } v = 0 \leq 0$ .

## Using Induction on $\Pi$ in Example

- **(B  $\rightarrow$  0BB)** Suppose  $w_1, w_2 \in \Pi_B$ , and assume the inductive hypothesis,  $w_1, w_2 \in Y$ . Thus  $w_1$  and  $w_2$  are nonempty. We must show that  $0w_1w_2 \in Y$ . Clearly,  $\text{diff}(0w_1w_2) = -1 + 1 + 1 = 1$ . So, suppose  $v$  is a proper prefix of  $0w_1w_2$ . We must show that  $\text{diff } v \leq 0$ . There are three cases to consider.
  - Suppose  $v = \%$ . Then  $\text{diff } v = 0 \leq 0$ .
  - Suppose  $v = 0u$ , for a proper prefix  $u$  of  $w_1$ . Because  $w_1 \in Y$ , we have that  $\text{diff } u \leq 0$ . Thus  $\text{diff } v = -1 + \text{diff } u \leq -1 + 0 \leq 0$ .

## Using Induction on $\Pi$ in Example

- **(B  $\rightarrow$  0BB)** Suppose  $w_1, w_2 \in \Pi_B$ , and assume the inductive hypothesis,  $w_1, w_2 \in Y$ . Thus  $w_1$  and  $w_2$  are nonempty. We must show that  $0w_1w_2 \in Y$ . Clearly,  $\text{diff}(0w_1w_2) = -1 + 1 + 1 = 1$ . So, suppose  $v$  is a proper prefix of  $0w_1w_2$ . We must show that  $\text{diff } v \leq 0$ . There are three cases to consider.
  - Suppose  $v = \%$ . Then  $\text{diff } v = 0 \leq 0$ .
  - Suppose  $v = 0u$ , for a proper prefix  $u$  of  $w_1$ . Because  $w_1 \in Y$ , we have that  $\text{diff } u \leq 0$ . Thus  $\text{diff } v = -1 + \text{diff } u \leq -1 + 0 \leq 0$ .
  - Suppose  $v = 0w_1u$ , for a proper prefix  $u$  of  $w_2$ . Because  $w_2 \in Y$ , we have that  $\text{diff } u \leq 0$ . Thus  $\text{diff } v = -1 + 1 + \text{diff } u = \text{diff } u \leq 0$ .

## Using Induction on $\Pi$ in Example

- **( $B \rightarrow 0BB$ )** Suppose  $w_1, w_2 \in \Pi_B$ , and assume the inductive hypothesis,  $w_1, w_2 \in Y$ . Thus  $w_1$  and  $w_2$  are nonempty. We must show that  $0w_1w_2 \in Y$ . Clearly,  $\text{diff}(0w_1w_2) = -1 + 1 + 1 = 1$ . So, suppose  $v$  is a proper prefix of  $0w_1w_2$ . We must show that  $\text{diff } v \leq 0$ . There are three cases to consider.
  - Suppose  $v = \%$ . Then  $\text{diff } v = 0 \leq 0$ .
  - Suppose  $v = 0u$ , for a proper prefix  $u$  of  $w_1$ . Because  $w_1 \in Y$ , we have that  $\text{diff } u \leq 0$ . Thus  $\text{diff } v = -1 + \text{diff } u \leq -1 + 0 \leq 0$ .
  - Suppose  $v = 0w_1u$ , for a proper prefix  $u$  of  $w_2$ . Because  $w_2 \in Y$ , we have that  $\text{diff } u \leq 0$ . Thus  $\text{diff } v = -1 + 1 + \text{diff } u = \text{diff } u \leq 0$ .
- The remaining productions are handled similarly.